



Subject Index to Volume 49

- Acellular collagen matrix
 augmented Achilles tendon repair in sheep, biomechanical evaluation of, 438–441
- Acellular porcine dermal patch
 cross-linked, and platelet-rich plasma fibrin matrix, for augmentation, comparison of Achilles tendon repair techniques in sheep model using, 128–134
- Achilles' tendon
 concomitant rupture of, and superior peroneal retinaculum (case report), 176–178
 insertion, variability of, 417–420
 lengthening of, for relief of forefoot pressure in patient with peripheral neuropathy, gastrocnemius recession as an alternative to (case report), 159.e9–e13
 reconstruction of, after sural fasciocutaneous flap using achilles tendon allograft with attached calcaneal bone block (case report), 86.e5–e10
 repair of
 in sheep, biomechanical evaluation of acellular collagen matrix augmented, 438–441
 in sheep model using cross-linked acellular porcine dermal patch and platelet-rich plasma fibrin matrix for augmentation, comparison of, 128–134
 uncomplicated method for minimally invasive, 208–211
 rupture of, reconstruction with combined V-Y plasty and gastrocnemius-soleus fascia turndown graft, 310–315
- Adenocarcinoma
 of colon, *Clostridium septicum* necrotizing fasciitis of forefoot secondary to (case report), 159.e1–e8
- Allograft
 Achilles tendon, with attached calcaneal bone block, achilles tendon reconstruction after sural fasciocutaneous flap using (case report), 86.e5–e10
 corticocancellous, treatment of unicameral bone cyst of calcaneus with endoscopic curettage and percutaneous filling with, 93–97
- American Society of Anesthesiologists
 in patients undergoing lower limb-preservation procedures, local anesthesia combined with monitored intravenous sedation for, 152–154
- Amputation
 transmetatarsal, preoperative Doppler assessment for, 101–105
- Anesthesia
 local
 combined with monitored intravenous sedation for American Society of Anesthesiologists in patients undergoing lower limb-preservation procedures, 152–154
 partial digital, randomized controlled trial to compare two techniques for, 143–146
- Aneurysmal bone cyst
 secondary, calcaneal chondroblastoma with (case report), 298.e5–e8
- Angioleiomyoma
 in ankle mimicking tarsal tunnel syndrome (case report), 398.e9–e15
- Ankle
 angioleiomyoma in, mimicking tarsal tunnel syndrome (case report), 398.e9–e15
 arthrodesis, outcome of, using a transfibular approach, 508–512
 arthroscopic treatment of anterolateral impingement of, 219–223
 arthroscopy of, delayed rupture of extensor hallucis longus and extensor digitorum communis tendons after breaching anterior capsule with radiofrequency probe during (case report), 490.e1–e3
 asymptomatic synovial chondromatosis of (case report), 565.e13–e17 and foot
 operative treatment for ganglion cysts of, 442–445
 reconstruction of, porcine SIS patch in, 123–127
 role of polymethylmethacrylate antibiotic-loaded cement in addition to debridement for treatment of soft tissue and osseous infections of, 55–62
- fractures of
 acute, factors associated with prolonged inpatient stay after surgical fixation of, 259–262
 missed posterior deep, inferior compartment syndrome in patient with (case report), 398.e5–e8
 irreducible fracture dislocation of, caused by tibialis posterior tendon interposition (case report), 166–171
 lateral, ligament of, new technique for stabilization of, 316 (Letter/Reply)
 in patients with diabetes and intact skin *versus* patients with diabetes and ulceration, bacterial skin contamination before and after surgical preparation of, 348–356
 percutaneous drilling for treatment of atraumatic osteonecrosis of, 20–24
 range of motion of, and functional outcome following total ankle arthroplasty, analysis of, 147–151
 surgically repaired bimalleolar fractures of, bimalleolar, use of noninvasive interactive neurostimulation to improve short-term recovery in patients with, 432–437
 transfibular arthrodesis, with fibular-onlay strut graft, 566–570
 use of negative pressure wound therapy for random local flaps at, 513–516
 use of Taylor Spatial Frame in compression arthrodesis of (case report), 182–187
- Ankle arthroplasty
 total, use of femoral distractor to aid distal tibial resection during, 205–207

- Ankylosis**
 tarsal, massive bilateral, and surgical correction (case report), 399.e11–e14
- Anterolateral impingement**
 of ankle, arthroscopic treatment of, 219–223
- Antibiotics**
 polymethylmethacrylate loaded cement with, role in addition to debridement for treatment of soft tissue and osseous infections of foot and ankle, 55–62
- Arthrodesis**
 ankle
 open, *versus* arthroscopic, 242–247
 outcome of, using a transfibular approach, 508–512
 calcaneocuboid distraction, *vs.* calcaneus osteotomy, results of lateral column lengthening by, 380–384
 chevron, of interphalangeal joint for hammertoe correction, 194–196
 compression, of ankle, use of Taylor Spatial Frame in (case report), 182–187
 distraction, of subtalar joint using allogeneic bone graft, 369–374
 Lapidus, modified, early weight bearing after, 357–362
 or total joint replacement, of the first metatarsophalangeal joint, 546–552
 primary, radiographic outcomes of first metatarsophalangeal joint in hallux abductovalgus deformity, 446–451
 subtalar and talonavicular, isolated medial incisional approach to, 326–330
 subtalar joint, using parallel guide and double screw fixation, 305–309
 of talonavicular joint, for flexible pes valgus deformity, 113–118
 tibiotalar, autogenous bone graft harvest using RIA technique for, 571–574
 transfibular ankle, with fibular-onlay strut graft, 566–570
- Arthroplasty**
 isolated Valenti, for hallux rigidus, need for surgical revision after, a systematic review, 294–297
 soft tissue interpositional, autogenous, for end-stage hallux rigidus, outcomes following, a systematic review, 475–478
 total ankle, analysis of ankle range of motion and functional outcome following, 147–151
- Arthroscopy**
 of ankle, delayed rupture of extensor hallucis longus and extensor digitorum communis tendons after breaching anterior capsule with radiofrequency probe during (case report), 490.e1–e3
versus open ankle arthrodesis, 242–247
 removal of intraarticular osteoid osteoma of distal tibia (case report), 398.e17–e21
 treatment of anterolateral impingement of ankle, 219–223
- Athletes**
 peroneal subluxation surgical results in, 238–241
- Bacterial skin**
 contamination before and after surgical preparation of foot, ankle, and lower leg in patients with diabetes and intact skin *versus* patients with diabetes and ulceration, 348–356
- Basal cell carcinoma**
 in diabetic patient, malignant transformation of chronic venous stasis ulcer to, 75–79
- Bed rest**
 effectiveness of instituting a specific bed protocol in reducing complications associated with, 340–347
- Biomechanics**
 evaluation of acellular collagen matrix augmented Achilles tendon repair in sheep, 438–441
- Blast injury**
 reconstruction of massive midfoot bone and soft tissue loss as result of, 301–304
- Bolster dressing**
 split-thickness skin graft, sterile cast padding as an alternative to commercially available cotton balls for, 98–100
- Bone cyst**
 aneurysmal, secondary, calcaneal chondroblastoma with (case report), 298.e5–e8
 unicameral, of calcaneus with endoscopic curettage and percutaneous filling with corticocancellous allograft, 93–97
- Bone graft**
 allogenic
 distraction arthrodesis of subtalar joint using, 369–374
 retrospective analysis of anterior calcaneal osteotomy with, 375–379
 autogenous, harvest of, using RIA technique for tibiotalar calcaneal arthrodesis, 571–574
 composite, substitute plugs, in treatment of chondral and osteochondral lesions of talus, effectiveness of, 224–231
- Book reviews**
 Zgonis: *Surgical Reconstruction of the Diabetic Foot and Ankle*, 107
- Brachymetatarsia**
 percutaneous distraction osteogenesis for treatment of, 197–204
- Button fixation**
 and tension wire, bilateral second metatarsal stress fractures after hallux valgus correction with use of (case report), 488.e15–e19
- Calcaneocuboid**
 distraction arthrodesis, *vs.* calcaneus osteotomy, results of lateral column lengthening by, 380–384
- Calcaneus**
 anterior osteotomy, with allogenic bone graft, retrospective analysis of, 375–379
 attached bone block, Achilles tendon reconstruction of after sural fasciocutaneous flap using achilles tendon allograft with (case report), 86.e5–e10
 chondroblastoma, with secondary aneurysmal bone cyst (case report), 298.e5–e8
 fractures of
 displaced intra-articular, in elderly patients, operative versus nonoperative treatment of, 25–32
 intra-articular, treatment with severe soft tissue damage hinged external fixator or internal stabilization, 8–15
 osteotomy of
 Evans, using anterior lumbar distractor-inserter, 401–403
vs. calcaneocuboid distraction arthrodesis, results of lateral column lengthening by, 380–384
 and plantar compartment, of foot, synovial sarcoma involving (case report), 398.e1–e4
 rupture of, acute, associated with ipsilateral malleolar fracture (case report), 565.e1–e4
 treatment of metastatic prostate adenocarcinoma to (case report), 159.e15–e20
 unicameral bone cyst of, treatment with endoscopic curettage and percutaneous filling with corticocancellous allograft, 93–97
- Case reports**
 Achilles tendon reconstruction after sural fasciocutaneous flap using Achilles tendon allograft with attached calcaneal bone block, 86.e5–e10

- acute calcaneus tendon rupture associated with ipsilateral malleolar fracture, 565.e1–e4
- angioleiomyoma in the ankle mimicking tarsal tunnel syndrome: a case report and review of the literature, 398.e9–e15
- an unusual etiology for adult-acquired flatfoot, 488.e11–488.e14
- application of an interosseous suture and button device for hallux valgus correction: a review of outcomes in a small series, 159.e21–e26
- arthroscopic removal of an intraarticular osteoid osteoma of the distal tibia, 398.e17–e21
- asymptomatic synovial chondromatosis of the ankle, 565.e13–e.17
- bilateral second metatarsal stress fractures after hallux valgus correction with the use of a tension wire and button fixation system, 488.e15–e19
- calcaneal chondroblastoma with secondary aneurysmal bone cyst, 298.e5–e8
- closed tibial fracture associated with laceration of tibialis anterior tendon, 86.e19–e22
- Clostridium septicum* necrotizing fasciitis of the forefoot secondary to adenocarcinoma of the colon, 159.e1–e8
- concomitant rupture of Achilles tendon and superior peroneal retinaculum, 176–178
- delayed rupture of the extensor hallucis longus and extensor digitorum communis tendons after breaching the anterior capsule with a radiofrequency probe during ankle arthroscopy, 490.e1–e3
- digital fracture after a flexor tendon transfer for hammertoe repair, 179–181
- disseminated intravascular coagulation, meningococcal infection, and ischemic changes affecting the lower extremities, 489.e5–e9
- gastrocnemius recession as an alternative to tendoAchillis lengthening for relief of forefoot pressure in a patient with peripheral neuropathy: a case report and description of a technical modification, 159.e9–e13
- giant solitary osteochondroma arising from the fifth metatarsal bone, 298.e9–e15
- high-pressure water injection of the foot with associated subcutaneous emphysema, 399.e15–e20
- hourglass ganglion cyst of the foot, 489.e11–e12
- intraaneural ganglion of the superficial peroneal nerve, 400.e1–e4
- irreducible dorsal metatarsophalangeal joint dislocation of the fifth toe, 298.e17–e20
- irreducible fracture dislocation of the ankle caused by tibialis posterior tendon interposition, 166–171
- irreducible Salter Harris type II distal tibial physeal fracture secondary to interposition of the posterior tibial tendon, 399.e5–e9
- late hematogenous infection of first metatarsophalangeal joint replacement: a case presentation, 489.e1–e4
- low-grade fibromyxoid sarcoma of the talus, 400.e5–e8
- Maisonneuve fracture without deltoid ligament disruption: a rare pattern of injury, 86.e11–e17
- massive bilateral tarsal ankylosis and surgical correction: a case report, 399.e11–e14
- melorheostosis: a literature review and case report with surgical considerations, 80–85
- missed posterior deep, inferior subcompartment syndrome in a patient with an ankle fracture, 398.e5–e8
- multicentric GCT of tarsal bones in an immature skeleton: a case report with review of literature, 399.e1–e4
- multifocal cementoplasty and internal fixation as a palliative procedure for skeletal metastases in the foot, 489.e13–e15
- myxoinflammatory fibroblastic sarcoma: a case report and review of the literature, 86.e1–e3
- periosteal chondrosarcoma of the foot, 488.e1–e4
- reimplantation of a totally extruded talus, 172–175
- retained hawthorn fragment in a child's foot complicated by infection: diagnosis and excision aided by localization with ultrasound, 161–165
- soft tissue anaplastic large T-cell lymphoma associated with a metallic orthopedic implant, 561–564
- stenosing tenosynovitis of the peroneus brevis tendon associated with hypertrophy of the peroneal tubercle, 188–190
- subcutaneous polypoid lipoma of the first intermetatarsal space mimicking rudimentary polydactyly, 298.e1–e4
- surgical repair of abductor hallucis muscle herniation, 488.e5–e9
- synovial sarcoma involving the calcaneus and plantar compartment of the foot, 398.e1–e4
- Tibialis posterior tendon rupture associated with a closed medial malleolar fracture, 565.e9–e12
- treatment of metastatic prostate adenocarcinoma to the calcaneus, 159.e15–e20
- the use of collagen injections in the treatment of metatarsalgia, 565.e5–e7
- use of the Taylor Spatial Frame in compression arthrodesis of the ankle, 182–187
- Cementoplasty
multifocal, and internal fixation, as palliative procedure for skeletal metastases of foot (case report), 489.e13–e15
- Charcot foot deformity
diabetic, reconstruction of, a two-stage percutaneous approach to, 517–522
- Cheilectomy
dorsal, for hallux rigidus using patient-reported outcome score, prospective assessment of, 232–237
isolated, for hallux rigidus, need for surgical revision after, a systematic review, 465–470
with phalangeal dorsiflexory osteotomy for hallux rigidus, outcomes after, a systematic review, 479–487
- Chondral lesions
of talus, effectiveness of composite bone graft substitute plugs in treatment of, 224–231
- Chondroblastoma
calcaneus, with secondary aneurysmal bone cyst (case report), 298.e5–e8
- Chondromatosis
asymptomatic synovial, of the ankle (case report), 565.e13–e.17
- Chondrosarcoma
periosteal, of foot (case report), 488.e1–e4
- Chopart joint injury
outcome and morbidity of, 541–545
- Clinical research
conducting, and getting the report published (Editorial), 109–111
human subject protection (Editorial), 317–318
- Clostridium septicum*
necrotizing fasciitis of forefoot secondary to adenocarcinoma of colon (case report), 159.e1–e8
- Clubfoot
comparison between Ponseti's and Kite's as treatment methods for, 395–397
treatment with Ponseti method, 426–431
- Collagen
injections of, use in the treatment of metatarsalgia (case report), 565.e5–e7

- Colon adenocarcinoma
Clostridium septicum necrotizing fasciitis of forefoot secondary to (case report), 159.e1–e8
- Complex regional pain syndrome (CRPS)
of lower extremity, long-term outcome after surgical treatment of, 33–36
- Computed tomography
dual isotope, optimized evaluation of diabetic foot infection by, 529–536
- Cotton balls
for split-thickness skin graft bolster dressing, sterile cast padding as an alternative to commercially available, 98–100
- Debridement
for treatment of soft tissue and osseous infections of foot and ankle, role of polymethylmethacrylate antibiotic-loaded cement in addition to, 55–62
- Deep venous thromboembolism
associated with bed rest, effectiveness of instituting a specific bed protocol in reducing, 340–347
- Deltoid ligament injury
Maisonneuve fractures without (case report), 86.e11–e17
- Diabetes mellitus
complications associated with uni-portal endoscopic gastrocnemius recession in patient with, 68–70
hallux ulceration in patients with, 2–7
and intact skin *versus* ulceration, bacterial skin contamination before and after surgical preparation of foot, ankle, and lower leg in patients with, 348–356
limb preservation, 106–107
limb-threatening and life-threatening in, 43–46
malignant transformation of chronic venous stasis ulcer to basal cell carcinoma in, 75–79
- Diabetic foot
charcot deformity, reconstruction of, a two-stage percutaneous approach to, 517–522
infection of, optimized evaluation by dual isotope SPECT/CT imaging protocol, 529–536
- Dislocation
of irreducible dorsal metatarsophalangeal joint, of fifth toe, 298.e17–e20
irreducible fracture, of ankle, caused by tibialis posterior tendon interposition (case report), 166–171
- Disseminated intravascular coagulation
changes affecting lower extremities (case report), 489.e5–e9
- Doppler assessment
preoperative, for transmetatarsal amputation, 101–105
- Drilling
percutaneous, for treatment of atraumatic osteonecrosis of ankle, 20–24
- Editorials
An Invitation to Authors to Submit Video Footage with Manuscripts, 111
Clarification, 112
Conducting Clinical Research and Getting the Report Published, 109–111
Health Care Reform, Statistical Significance, Effect Size, and the Future of the Profession, 319–320
Human Subject Protection: Overkill?, 317–318
A New Look for JFAS, 1
Successful Treatment Hinges on an Accurate Diagnosis, 1
- “Take Your Government Hands Off My Medicare”, 409–410
Time Well Spent, 217–218
Who Is Packing Your Parachute?, 507
- Elderly
displaced intra-articular calcaneal fracture, operative versus nonoperative treatment in, 25–32
- Emphysema
subcutaneous, high-pressure water injection of foot associated with (case report), 399.e15–e20
- Endoscopic curettage
and percutaneous filling, with corticocancellous allograft, treatment of unicameral bone cyst of calcaneus with, 93–97
- Equine pericardium
xenograft, implanted, inflammatory reaction to, 155–158
- Extensor digitorum communis tendons
delayed rupture of, after breaching anterior capsule with a radiofrequency probe during ankle arthroscopy (case report), 490.e1–e3
- Extensor hallucis longus
delayed rupture of, after breaching anterior capsule with a radiofrequency probe during ankle arthroscopy (case report), 490.e1–e3
- External fixation
circular, assembly of, kickstand technique for, 504–505
pin sites, prevention and management of complications arising from, 87–92
- Femoral distractor
use to aid distal tibial resection during total ankle arthroplasty, 205–207
- Fibromyxoid sarcoma
low-grade, of talus (case report), 400.e5–e8
- Fibular fracture
high, in association with triplane fracture, 491–494
- Fixation
surgical, of acute ankle fracture, factors associated with prolonged inpatient stay after, 259–262
- Flaps
local, at the ankle region, use of negative pressure wound therapy for, 513–516
sural fasciocutaneous, using Achilles tendon allograft with attached calcaneal bone block, Achilles tendon reconstruction after (case report), 86.e5–e10
- Flatfoot
adult, in United States, characteristics of, 363–368
adult-acquired, an unusual etiology for (case report), 488.e11–488.e14
- Flexible pes valgus deformity
talonavicular joint arthrodesis for, 113–118
- Flexor digitorum longus
transfer for posterior tibial tendon dysfunction, interference screw fixation and short harvest using, 501–503
- Flexor tendon transfer
for hammertoe repair, digital fracture after (case report), 179–181
- Foot
and ankle
operative treatment for ganglion cysts of, 442–445
reconstruction of, porcine SIS patch in, 123–127
role of polymethylmethacrylate antibiotic-loaded cement in addition to debridement for treatment of soft tissue and osseous infections of, 55–62
with associated subcutaneous emphysema, high-pressure water injection of (case report), 399.e15–e20

- Charcot diabetic, a two-stage percutaneous approach for reconstruction of, 517–522
- in child, complicated by infection, retained hawthorn fragment in (case report), 161–165
- hourglass ganglion cyst of (case report), 489.e11–e12
- multifocal cementoplasty and internal fixation as palliative procedure for skeletal metastases in (case report), 489.e13–e15
- in patients with diabetes and intact skin *versus* patients with diabetes and ulceration, bacterial skin contamination before and after surgical preparation of, 348–356
- periosteal chondrosarcoma of (case report), 488.e1–e4
- primary nodular melanoma of, 263–273
- soft tissue defects of, in patient at high risk, V-Y fasciocutaneous advancement flap coverage of, 71–74
- synovial sarcoma involving the calcaneus and plantar compartment of, 398.e1–e4
- Forefoot**
- pressure of, in patient with peripheral neuropathy, gastrocnemius recession as an alternative to tendoAchillis lengthening for relief of (case report), 159.e9–e13
- secondary to adenocarcinoma of colon, *Clostridium septicum* necrotizing fasciitis of (case report), 159.e1–e8
- Fracture**
- of ankle
- acute, factors associated with prolonged inpatient stay after surgical fixation of, 259–262
 - irreducible dislocation of, caused by tibialis posterior tendon interposition (case report), 166–171
 - missed posterior deep, inferior subcompartment syndrome in patient with (case report), 398.e5–e8
 - surgically repaired bimalleolar bimalleolar, use of noninvasive interactive neurostimulation to improve short-term recovery in patients with, 432–437
- of bilateral second metatarsal stress, after hallux valgus correction with the use of tension wire and button fixation system (case report), 488.e15–e19
- calcaneal
- displaced intra-articular, in elderly patients, operative versus nonoperative treatment of, 25–32
 - intra-articular, treatment with severe soft tissue damage hinged external fixator or internal stabilization, 8–15
- closed tibial, associated with laceration of tibialis anterior tendon (case report), 86.e19–e22
- digital, after flexor tendon transfer for hammertoe repair (case report), 179–181
- of Maisonneuve
- successful treatment hinges on an accurate diagnosis (Editorial), 1
 - without deltoid ligament disruption (case report), 86.e11–e17
- of medial malleolar, comparison of pullout strength between 3.5-mm bicortical screws vs. 4.0-mm partially threaded cancellous screws in fixation of, 248–252
- of pilon, short-term financial outcomes of, 47–51
- of Salter Harris type II distal tibial physeal, secondary to interposition of posterior tibial tendon (case report), 399.e5–e9
- triplane, high fibular fracture in association with, 491–494
- Fusion.** *see* Arthrodesis
- Ganglion**
- intra-neural
- of the superficial peroneal nerve, 579 (Letters)
 - of superficial peroneal nerve (case report), 400.e1–e4
- Ganglion cyst
- of foot and ankle, operative treatment for, 442–445
 - hourglass, of foot (case report), 489.e11–e12
 - superficial peroneal intraneural, originating from the inferior tibiofibular joint, 575–578
- Gastrocnemius recession**
- as an alternative to tendoAchillis lengthening for relief of forefoot pressure in patient with peripheral neuropathy (case report), 159.e9–e13
 - extended knee hemilithotomy position for, 214–216
 - uni-portal endoscopic, in diabetic patient population, complications associated with, 68–70
- Gastrocnemius-soleus fascia turndown graft and V-Y plasty, reconstruction of Achilles tendon rupture with combined, 310–315**
- Giant cell tumor**
- multicentric, of tarsal bones in immature skeleton (case report), 399.e1–e4
- Grafts**
- fibular-onlay strut, transfibular ankle arthrodesis with, 566–570
 - gastrocnemius-soleus fascia turndown, combined with V-Y plasty, reconstruction of Achilles tendon rupture with combined, 310–315
 - skin, full- and split-thickness, VERSAJET™ hydrosurgery technique for, 404–407
- Hallux abductovalgus**
- deformity of, radiographic outcomes following primary arthrodesis of first metatarsophalangeal joint in, 446–451
- Hallux rigidus**
- clinical outcomes after isolated periarticular osteotomies of first metatarsal for, a systematic review, 553–560
 - end-stage, outcomes following autogenous soft tissue interpositional arthroplasty for, a systematic review, 475–478
 - need for surgical revision after isolated cheilectomy for, a systematic review, 465–470
 - need for surgical revision after isolated Valenti arthroplasty for, a systematic review, 294–297
 - outcomes after cheilectomy with phalangeal dorsiflexory osteotomy for, a systematic review, 479–487
 - surgical treatment using metatarsal head resurfacing implant, 321–325
 - using patient-reported outcome score, prospective assessment of dorsal cheilectomy for, 232–237
- Hallux ulceration**
- in diabetic patients, 2–7
- Hallux valgus**
- correction of, application of an interosseous suture and button device for (case report), 159.e21–e26
 - correction with the use of tension wire and button fixation system, bilateral second metatarsal stress fractures after (case report), 488.e15–e19
 - radiographic angles in, comparison between manual and computer-assisted measurements for, 523–528
 - scarf-Akin osteotomy correction for, 16–19
- Hammertoe**
- correction of, chevron arthrodesis of interphalangeal joint for, 194–196
 - repair of, digital fracture after flexor tendon transfer for (case report), 179–181
- Hawthorn**
- retained fragment of, in child's foot complicated by infection (case report), 161–165

- Health care reform
 statistical significance, effect size, and the future of the profession
 (Editorial), 319–320
- Heel
 pain, diagnosis and treatment of, S1–S19
- Hemiarthroplasty
 BioPro® prosthesis, retrospective cohort study of, 331–339
- Hemilithotomy
 extended knee position, for gastrocnemius recession, 214–216
- Herniation
 of abductor hallucis muscle, surgical repair of (case report),
 488.e5–.e9
- Hinged external fixator
 treatment of intra-articular calcaneus fractures with severe soft
 tissue damage with, 8–15
- Hydrosurgery
 VERSAJET™, technique for full- and split-thickness skin graft, 404–407
- Hypertrophy
 of peroneal tubercle, stenosing tenosynovitis of peroneus brevis
 tendon associated with (case report), 188–190
- Implant
 equine pericardium xenograft, inflammatory reaction to, 155–158
 metallic orthopedic, soft tissue anaplastic large T-cell lymphoma
 associated with (case report), 561–564
 metatarsal head resurfacing, surgical treatment of hallux rigidus
 using, 321–325
- Infections
 of diabetic foot, optimized evaluation by dual isotope SPECT/CT
 imaging protocol, 529–536
 late hematogenous, of first metatarsophalangeal joint replacement
 (case report), 489.e1–e4
 lower extremity wound, efficacy of power-pulsed lavage in, 135–142
 meningococcal, changes affecting lower extremities (case report),
 489.e5–e9
 retained hawthorn fragment in a child's foot complicated by (case
 report), 161–165
 soft tissue and osseous, of foot and ankle, role of
 polymethylmethacrylate antibiotic-loaded cement in
 addition to debridement for treatment of, 55–62
- Injections
 of collagen, use in the treatment of metatarsalgia (case report),
 565.e5–e7
- Injury
 of Chopart joint, outcome and morbidity of, 541–545
- Intermetatarsal space
 first, mimicking rudimentary polydactyly, subcutaneous polypoid
 lipoma of (case report), 298.e1–e4
- Internal fixation
 and multifocal cementoplasty, as palliative procedure for skeletal
 metastases of foot (case report), 489.e13–e15
- Internal stabilization
 treatment of intra-articular calcaneus fractures with severe soft
 tissue damage with, 8–15
- Interphalangeal joint
 for hammertoe correction, chevron arthrodesis of, 194–196
- Ischemia
 changes affecting lower extremities (case report), 489.e5–e9
- Joint replacement
 total, or arthrodesis, of the first metatarsophalangeal joint,
 546–552
- Journal of Foot and Ankle Surgery*
 publication of manuscripts in (Editorial), 1
 5-year review of statistical methods presented in, 471–474
- Kickstand technique
 for circular external fixator assembly, 504–505
- Kidner-Cobb procedure
 modified, for symptomatic flexible pes planovalgus and posterior
 tibial tendon dysfunction stage II, 411–416
- Kite method
 vs. Ponseti's, for treatment of clubfoot, 395–397
- Knee position
 extended, hemilithotomy, for gastrocnemius recession, 214–216
- Laceration
 of tibialis anterior tendon, closed tibial fracture associated with
 (case report), 86.e19–e22
- Lapidus arthrodesis
 modified, immediate weight bearing following, 459–464
- Lateral ligament, ankle
 new technique for stabilization of, 316 (Letter/Reply)
- Lengthening procedures
 results of lateral column, by calcaneus osteotomy vs.
 calcaneocuboid distraction arthrodesis, 380–384
 of tendoAchillis, for relief of forefoot pressure in patient with
 peripheral neuropathy, gastrocnemius recession as an
 alternative to (case report), 159.e9–e13
- Life-threatening
 and limb-threatening, diabetic extremities, 43–46
- Limb preservation
 diabetic, 106–107
- Limb salvage
 safety of local anesthesia combined with monitored intravenous
 sedation for American Society of Anesthesiologists in
 patients undergoing, 152–154
- Limb-threatening
 and life-threatening, diabetic extremities, 43–46
- Local anesthesia
 combined with monitored intravenous sedation for American
 Society of Anesthesiologists in patients undergoing lower
 limb-preservation procedures, 152–154
 partial digital, randomized controlled trial to compare two
 techniques for, 143–146
- Lower extremity
 changes experienced during pregnancy, 452–458
 CRPS, long-term outcome after surgical treatment of, 33–36
 disseminated intravascular coagulation, meningococcal infection,
 and ischemic changes affecting (case report), 489.e5–e9
 wound infection of, efficacy of power-pulsed lavage in, 135–142
- Lower leg
 in patients with diabetes and intact skin *versus* patients with
 diabetes and ulceration, bacterial skin contamination before
 and after surgical preparation of, 348–356
- Lumbar distractor-inserter
 anterior, Evans calcaneal osteotomy using, 401–403
- Lymphoma
 of T-cell, soft tissue anaplastic, associated with a metallic
 orthopedic implant (case report), 561–564
- Maisonneuve
 fractures of, successful treatment hinges on an accurate diagnosis
 (Editorial), 1

- Malleolar fracture
 closed medial, tibialis posterior tendon rupture associated with (case report), 565.e9–e12
 ipsilateral, acute calcaneus tendon rupture associated with (case report), 565.e1–e4
 medial, comparison of pullout strength between 3.5-mm bicortical screws vs. 4.0-mm partially threaded cancellous screws in fixation of, 248–252
- Medicare
 government-run health care model (Editorial), 409–410
- Melanoma
 primary nodular, of foot, 263–273
- Melorheostosis
 with surgical intervention (case report), 80–85
- Meningococcal infection
 changes affecting lower extremities (case report), 489.e5–e9
- Metastasis
 of prostate adenocarcinoma to the calcaneus, 159.e15–e20
 skeletal, of foot, multifocal cementoplasty and internal fixation as palliative procedure for (case report), 489.e13–e15
- Metatarsal bones
 fifth, giant solitary osteochondroma arising from (case report), 298.e9–e15
 Scarf and Weil osteotomy, of lateral rays for correction of rheumatoid forefoot deformities, a systematic review, 390–394
- Metatarsalgia
 use of collagen injections in the treatment of (case report), 565.e5–e7
- Metatarsal head
 resurfacing implant, surgical treatment of hallux rigidus using, 321–325
- Metatarsal joint
 first, for hallux rigidus, clinical outcomes after isolated periarticular osteotomies of, a systematic review, 553–560
- Metatarsal osteotomy
 first, multiplanar effect on sesamoid position, 63–67
- Metatarsal ray
 resection of, minimum-incision in, 52–54
- Metatarsal stress
 bilateral second, fractures of, after hallux valgus correction with the use of tension wire and button fixation system (case report), 488.e15–e19
- Metatarsophalangeal joint
 first
 in hallux abductovalgus deformity, radiographic outcomes of primary arthrodesis of, 446–451
 replacement of, late hematogenous infection of (case report), 489.e1–e4
 replacement of, using Moje, 37–42
 total joint replacement or arthrodesis of, 546–552
 unusual injury to, 212–213
 irreducible dorsal dislocation of, in fifth toe (case report), 298.e17–e20
- Midfoot bone
 massive, and soft tissue loss, as result of blast injury, reconstruction of, 301–304
- Minimal incision surgery
 in metatarsal ray resection, 52–54
- Minimally invasive surgery
 of Achilles tendon repair, uncomplicated method for, 208–211
- Moje
 first metatarsophalangeal joint replacement using, 37–42
- Musculotendinous junction
 abnormal, proximal, of peroneus brevis muscle as cause of peroneus brevis tendon tears, 537–540
 Myxoinflammatory fibroblastic sarcoma (case report), 86.e1–e3
- Nail
 puncture wound, through rubber-soled shoe, 421–425
- Necrotizing fasciitis
Clostridium septicum, of forefoot secondary to adenocarcinoma of colon (case report), 159.e1–e8
- Negative pressure wound therapy
 use for random local flaps at the ankle region, 513–516
- Neuropathy
 peripheral, gastrocnemius recession as an alternative to tendoAchillis lengthening for relief of forefoot pressure in patient with (case report), 159.e9–e13
- Neurostimulation
 noninvasive interactive, use to improve short-term recovery in patients with surgically repaired bimalleolar ankle fractures, 432–437
- Osseous infection
 of foot and ankle, role of polymethylmethacrylate antibiotic-loaded cement in addition to debridement for treatment of, 55–62
- Osteochondral lesions
 of talus, effectiveness of composite bone graft substitute plugs in treatment of, 224–231
- Osteochondroma
 giant solitary, arising from fifth metatarsal bone (case report), 298.e9–e15
- Osteogenesis
 percutaneous distraction, for treatment of brachymetatarsia, 197–204
- Osteoid osteoma
 intraarticular, of distal tibia, arthroscopy removal of (case report), 398.e17–e21
- Osteonecrosis
 atraumatic, of ankle, percutaneous drilling for treatment of, 20–24
- Osteotomy
 calcaneal
 anterior, with allogenic bone graft, retrospective analysis of, 375–379
 Evans, using anterior lumbar distractor-inserter, 401–403
 calcaneus, vs. calcaneocuboid distraction arthrodesis, results of lateral column lengthening by, 380–384
 isolated periarticular, of first metatarsal for hallux rigidus, clinical outcomes after, a systematic review, 553–560
 metatarsal, first, multiplanar effect on sesamoid position, 63–67
 phalangeal dorsiflexory, for hallux rigidus, outcomes after cheilectomy with, a systematic review, 479–487
 scarf-Akin, for hallux valgus, 16–19
 Scarf and Weil metatarsal, of lateral rays for correction of rheumatoid forefoot deformities, a systematic review, 390–394
- Outcomes
 after cheilectomy with phalangeal dorsiflexory osteotomy for hallux rigidus, a systematic review, 479–487
 of ankle arthrodesis using transfibular approach, 508–512
 clinical, after isolated periarticular osteotomies of first metatarsal for hallux rigidus, a systematic review, 553–560
 following autogenous soft tissue interpositional arthroplasty for end-stage hallux rigidus, a systematic review, 475–478

- functional, following total ankle arthroplasty, analysis of ankle range of motion and, 147–151
- long-term, after surgical treatment of lower extremity CRPS, 33–36
- and morbidity of Chopart joint injury, 541–545
- radiographic, following primary arthrodesis of first metatarsophalangeal joint in hallux abducto valgus deformity, 446–451
- short-term financial, of pilon fractures, 47–51
- Pain
 - in heel, diagnosis and treatment of, S1–S19
- Patient-reported outcome score
 - prospective assessment of dorsal cheilectomy for hallux rigidus using, 232–237
- Percutaneous distraction
 - osteogenesis, for treatment of brachymetatarsia, 197–204
- Percutaneous filling
 - and endoscopic curettage, with corticocancellous allograft, treatment of unicameral bone cyst of calcaneus with, 93–97
- Periarticular osteotomy
 - isolated, of first metatarsal for hallux rigidus, clinical outcomes after, a systematic review, 553–560
- Periosteal chondrosarcoma
 - of foot (case report), 488.e1–e4
- Peripheral neuropathy
 - gastrocnemius recession as an alternative to tendoAchillis lengthening for relief of forefoot pressure in patient with (case report), 159.e9–e13
- Peroneal nerve
 - superficial
 - intra-neural ganglion of, 579 (Letters)
 - intra-neural ganglion of (case report), 400.e1–e4
- Peroneal retinaculum
 - superior, and concomitant rupture of Achilles tendon (case report), 176–178
- Peroneal subluxation
 - surgical results in athletic patients, 238–241
- Peroneal tubercle
 - stenosing tenosynovitis of peroneus brevis tendon associated with hypertrophy of (case report), 188–190
- Peroneus brevis muscle
 - as cause of peroneus brevis tendon tears, abnormal proximal musculotendinous junction of, 537–540
- Peroneus brevis tendon
 - associated with hypertrophy of peroneal tubercle, stenosing tenosynovitis of (case report), 188–190
- Peroneus longus tendon
 - split, in lateral ankle ligament stabilization, 316 (Letter/Reply)
- Pes planovalgus
 - symptomatic flexible, and posterior tibial tendon dysfunction stage II, modified Kidner-Cobb procedure for, 411–416
 - and tarsal coalition, review of, 274–293
- Phalangeal dorsiflexory
 - osteotomy of, for hallux rigidus, outcomes after cheilectomy with, a systematic review, 479–487
- Pilon fractures
 - short-term financial outcomes of, 47–51
- Pin sites
 - prevention and management of complications arising from external fixation, 87–92
- Plantar compartment
 - and calcaneus, of foot, synovial sarcoma involving (case report), 398.e1–e4
- Plantar ulcers
 - of toes, functional outcome and patient satisfaction after flexor tenotomy for, 119–122
- Platelet gel
 - autologous, use in toenail surgery, 385–389
- Platelet-rich plasma fibrin matrix
 - and cross-linked acellular porcine dermal patch, for augmentation, comparison of Achilles tendon repair techniques in sheep model using, 128–134
- Pneumonia
 - hospital-acquired, associated with bed rest, effectiveness of instituting a specific bed protocol in reducing, 340–347
- Polymethylmethacrylate
 - antibiotic-loaded cement, role in addition to debridement for treatment of soft tissue and osseous infections of foot and ankle, 55–62
- Polypoid lipoma
 - subcutaneous, of first intermetatarsal space mimicking rudimentary polydactyly (case report), 298.e1–e4
- Ponseti method
 - treatment of clubfoot with, 426–431
 - vs. Kite's, for treatment of clubfoot, 395–397
- Power-pulsed lavage
 - efficacy in lower extremity wound infections, 135–142
- Pregnancy
 - lower extremity changes experienced during, 452–458
- Pressure ulceration
 - associated with bed rest, effectiveness of instituting a specific bed protocol in reducing, 340–347
- Prostate adenocarcinoma
 - metastatic, to calcaneus, treatment of (case report), 159.e15–e20
- Prosthesis
 - retrospective cohort study of BioPro® hemiarthroplasty, 331–339
- Publications
 - of report, and conducting clinical research (Editorial), 109–111
- Pullout strength
 - comparison between 3.5-mm bicortical screws vs. 4.0-mm partially threaded cancellous screws in fixation of medial malleolar fractures, 248–252
- Radiofrequency probe
 - during ankle arthroscopy, delayed rupture of extensor hallucis longus and extensor digitorum communis tendons after breaching anterior capsule with (case report), 490.e1–e3
- Radiography
 - for hallux valgus, comparison between manual and computer-assisted measurements in, 523–528
 - outcomes following primary arthrodesis of first metatarsophalangeal joint in hallux abducto valgus deformity, 446–451
- Reamer Irrigator Aspirator technique
 - for tibiotalar calcaneal arthrodesis, autogenous bone graft harvest using, 571–574
- Reconstruction
 - of Achilles tendon, after sural fasciocutaneous flap using Achilles tendon allograft with attached calcaneal bone block (case report), 86.e5–e10
 - of Achilles tendon rupture with combined V-Y plasty and gastrocnemius-soleus fascia turndown graft, 310–315
 - of Charcot diabetic foot, a two-stage percutaneous approach to, 517–522
 - of foot and ankle, porcine SIS patch in, 123–127

- of massive midfoot bone and soft tissue loss as result of blast injury, 301–304
- Reimplantation
 - of totally extruded talus (case report), 172–175
- Rheumatoid forefoot deformity
 - correction of, Scarf and Weil metatarsal osteotomies of lateral rays for, a systematic review, 390–394
- RIA. *see* Reamer Irrigator Aspirator
- Rubber-soled shoe
 - nail puncture wound through, 421–425
- Rudimentary polydactyly
 - subcutaneous polypoid lipoma of first intermetatarsal space mimicking (case report), 298.e1–e4
- Rupture
 - of calcaneus tendon, acute, associated with ipsilateral malleolar fracture (case report), 565.e1–e4
 - of tibialis posterior tendon, associated with a closed medial malleolar fracture (case report), 565.e9–e12
- Salter Harris
 - type II, irreducible, distal tibial physeal fracture secondary to interposition of the posterior tibial tendon (case report), 399.e5–e9
- Sarcoma
 - fibromyxoid, low-grade, of talus (case report), 400.e5–e8
 - myxoinflammatory fibroblastic (case report), 86.e1–e3
 - synovial, involving calcaneus and plantar compartment of foot (case report), 398.e1–e4
- Screw fixation
 - double, subtalar joint fusion using parallel guide and, 305–309
 - interference, and short harvest, using flexor digitorum longus transfer for posterior tibial tendon dysfunction, 501–503
 - of medial malleolar fractures, comparison of pullout strength between 3.5-mm vs. bicortical screws and 4.0-mm partially threaded cancellous screw in, 248–252
- Sedation
 - intravenous, monitored, for American Society of Anesthesiologists in patients undergoing lower limb-preservation procedures, local anesthesia combined with, 152–154
- Sesamoid bone
 - position of, multiplanar effect of first metatarsal osteotomy on, 63–67
- Severe open ankle sprain (SOAS)
 - lesion presenting as penetrating soft tissue injury, 253–258
- Single-photon emission computed tomography
 - dual isotope, optimized evaluation of diabetic foot infection by, 529–536
- Skeleton
 - immature, multicentric GCT of tarsal bones in (case report), 399.e1–e4
- Skin graft
 - full- and split-thickness, VERSAJET™ hydrosurgery technique for, 404–407
- Small intestinal submucosa (SIS)
 - porcine patch, in foot and ankle reconstruction, 123–127
- Soft tissue damage
 - severe, with hinged external fixator or internal stabilization, treatment of intra-articular calcaneus fractures with, 8–15
- Soft tissue defects
 - of foot in patient at high risk, V-Y fasciocutaneous advancement flap coverage of, 71–74
- Soft tissue infection
 - of foot and ankle, role of polymethylmethacrylate antibiotic-loaded cement in addition to debridement for treatment of, 55–62
- Soft tissue injury
 - penetrating, SOAS lesion presenting as, 253–258
- Soft tissue loss
 - and massive midfoot bone, as result of blast injury, reconstruction of, 301–304
- Sterile cast padding
 - as an alternative to commercially available cotton balls for split-thickness skin graft bolster dressing, 98–100
- Subcompartment syndrome
 - inferior, missed posterior deep, in patient with ankle fracture (case report), 398.e5–e8
- Subtalar
 - arthrodesis, isolated medial incisional approach to, 326–330
- Subtalar joint
 - fusion, using parallel guide and double screw fixation, 305–309
 - using allogeneic bone graft, distraction arthrodesis of, 369–374
- Surgery
 - repair of abductor hallucis muscle herniation, 488.e5–.e9
 - toenail, use of autologous platelet gel in, 385–389
- Surgical incision
 - high-risk, negative-pressure wound therapy applied to, 299–300
- Suture
 - interosseous, and button device, application for hallux valgus correction (case report), 159.e21–e26
- Syndesmosis
 - tibiofibular, distal, tri-ligamentous reconstruction of, 495–500
- Synovial chondromatosis
 - asymptomatic, of the ankle (case report), 565.e13–e.17
- Synovial sarcoma
 - involving calcaneus and plantar compartment of foot (case report), 398.e1–e4
- Talonavicular
 - arthrodesis, isolated medial incisional approach to, 326–330
- Talonavicular joint
 - arthrodesis of, for flexible pes valgus deformity, 113–118
- Talus
 - effectiveness of composite bone graft substitute plugs in treatment of chondral and osteochondral lesions of, 224–231
 - low-grade fibromyxoid sarcoma of (case report), 400.e5–e8
 - totally extruded, reimplantation of (case report), 172–175
- Tarsal ankylosis
 - massive bilateral, and surgical correction (case report), 399.e11–e14
- Tarsal bones
 - in immature skeleton, multicentric GCT of (case report), 399.e1–e4
- Tarsal coalition
 - and pes planovalgus, review of, 274–293
- Tarsal tunnel syndrome
 - angioleiomyoma in ankle mimicking (case report), 398.e9–e15
- Taylor Spatial Frame (TSF)
 - use in compression arthrodesis of ankle (case report), 182–187
- T-cell lymphoma
 - large, soft tissue anaplastic, associated with a metallic orthopedic implant (case report), 561–564
- Tears
 - of peroneus brevis tendon, abnormal proximal musculotendinous junction of peroneus brevis muscle as cause of, 537–540
- Tendons
 - peroneus brevis, tears of, abnormal proximal musculotendinous junction of peroneus brevis muscle as cause of, 537–540
 - tibialis anterior, laceration of, closed tibial fracture associated with (case report), 86.e19–e22

- Tenosynovitis
 stenosing, of peroneus brevis tendon associated with hypertrophy of peroneal tubercle (case report), 188–190
- Tenotomy
 flexor, for plantar ulcers of toes, functional outcome and patient satisfaction after, 119–122
- Tension wire
 and button fixation, bilateral second metatarsal stress fractures after hallux valgus correction with use of (case report), 488.e15–e19
- Tibia
 distal
 arthroscopic removal of intraarticular osteoid osteoma of (case report), 398.e17–e21
 resection of, during total ankle arthroplasty, use of femoral distractor to aid, 205–207
- Tibial fracture
 closed, associated with laceration of tibialis anterior tendon (case report), 86.e19–e22
- Tibial tendon
 anterior, laceration of, closed tibial fracture associated with (case report), 86.e19–e22
 posterior
 dysfunction of, interference screw fixation and short harvest using flexor digitorum longus transfer for, 501–503
 dysfunction of, stage II, modified Kidner-Cobb procedure for symptomatic flexible pes planovalgus and, 411–416
 interposition, irreducible fracture dislocation of ankle caused by (case report), 166–171
 irreducible Salter Harris type II distal tibial physeal fracture secondary to interposition of (case report), 399.e5–e9
 rupture of, associated with closed medial malleolar fracture (case report), 565.e9–e12
- Tibiofibular joint
 inferior, superficial peroneal intraneural ganglion cyst originating from, 575–578
- Tibiofibular syndesmosis
 distal, tri-ligamentous reconstruction of, 495–500
- Toe
 fifth, irreducible dorsal metatarsophalangeal joint dislocation of (case report), 298.e17–e20
 functional outcome and patient satisfaction after flexor tenotomy for plantar ulcers of, 119–122
- Toenail
 surgery, use of autologous platelet gel in, 385–389
- Total joint replacement. *see* Joint replacement, total
- Transfibular approach
 outcome of ankle arthrodesis using, 508–512
- Transmetatarsal amputation
 preoperative Doppler assessment for, 101–105
- Tri-ligamentous
 reconstruction of, distal tibiofibular syndesmosis, 495–500
- Ulceration
 bacterial skin contamination before and after surgical preparation of foot, ankle, and lower leg in patients with diabetes and intact skin *versus* patients with, 348–356
 of hallux, in diabetic patients, 2–7
 plantar, of toe, functional outcome and patient satisfaction after flexor tenotomy for, 119–122
- Unicameral bone cyst
 of calcaneus with endoscopic curettage and percutaneous filling with corticocancellous allograft, 93–97
- United States
 characteristics of adult flatfoot in, 363–368
- Venous stasis ulcer
 chronic, to basal cell carcinoma, in diabetic patient, malignant transformation of, 75–79
- VERSAJET™
 hydrosurgery technique for full- and split-thickness skin graft, 404–407
- Video footage
 with manuscripts, an invitation to authors to submit (Editorial), 111
- V-Y fasciocutaneous advancement flap
 coverage of soft tissue defects of foot in patient at high risk, 71–74
- V-Y plasty
 combined with gastrocnemius-soleus fascia turndown graft, reconstruction of Achilles tendon rupture with combined, 310–315
- Water injection
 high-pressure, of foot with associated subcutaneous emphysema (case report), 399.e15–e20
- Weight bearing
 early, after modified Lapidus arthrodesis, 357–362
 immediate, following modified Lapidus arthrodesis, 459–464
- Wound
 infection, of lower extremity, efficacy of power-pulsed lavage in, 135–142
 nail puncture, through rubber-soled shoe, 421–425
- Wound therapy
 negative-pressure, applied to high-risk surgical incisions, 299–300
- Xenograft
 implanted equine pericardium, inflammatory reaction to, 155–158
- Z-plasty
 precise intraoperative 60° skin, 191–193